



A rare cause of deep peroneal nerve palsy due to compression of synovial cyst – Case report

Mehmet Erdil^{a,*}, Korhan Ozkan^b, Feyza Unlu Ozkan^c, Kerem Bilsel^a, Ismail Turkmen^b, Serkan Senol^d, Serhan Sarar^e

^a Bezmialem Vakif University, Medicine Faculty, Department of Orthopaedics and Traumatology, Istanbul, Turkey

^b Istanbul Medeniyet Universitesi, Faculty of Medicine, Department of Ortopaedics and Traumatology, Istanbul, Turkey

^c Fatih Sultan Mehmet Research and Training Hospital, Department of Physical Therapy and Rehabilitation, Istanbul, Turkey

^d Istanbul Medeniyet Universitesi, Faculty of Medicine, Department of Patology, Istanbul, Turkey

^e Istanbul Medeniyet Universitesi, Faculty of Medicine, Department of Anesthesia and Reanimation, Istanbul, Turkey

ARTICLE INFO

Article history:

Received 16 August 2012

Received in revised form

10 November 2012

Accepted 27 November 2012

Available online 13 March 2013

Keywords:

Synovial cyst

Peroneal nerve

Compression

Neuropathy

ABSTRACT

INTRODUCTION: Synovial cyst is a rare cause of compression neuropathy and its differential diagnosis can be misleading.

PRESENTATION OF CASE: This article presents clinical, radiological, and histological findings of deep peroneal nerve palsy due to compression of a synovial cyst in a 30-year-old patient admitted with sudden drop foot.

DISCUSSION: Focal nerve entrapment in lower extremity due to synovial cyst is a rare entity. Differential diagnosis is important. Surgical excision is the main treatment method with high success rate.

CONCLUSION: Synovial cyst compression which can be treated easily with surgical excision should be considered in rapidly progressed drop foot.

© 2013 Surgical Associates Ltd. Published by Elsevier Ltd. Open access under [CC BY-NC-ND license](http://creativecommons.org/licenses/by-nc-nd/4.0/).

1. Introduction

Although synovial cysts are very common soft tissue tumors, compression neuropathy due to synovial cysts are extremely rare. Several cases of ulnar and median nerve compressions due to ganglia have been described.^{1–3} However, synovial cysts result peripheral nerve compression in the lower extremities much more rarely. Sultan firstly reported a case of compression neuropathy to peroneal nerve by a synovial cyst in 1921.⁴

In the present case report, a rarely seen compression neuropathy of deep peroneal nerve by a synovial cyst has been presented and evaluated in the light of literature.

2. Presentation of case

A 30-year-old man was admitted to our hospital complaining of pain in lateral part of his proximal leg and drop foot that

progressed rapidly during the same day in several hours. On physical examination, there were no atrophy of the peroneus muscles and the tibialis anterior muscle, however, peroneal muscles strength was 5/5 whereas tibialis anterior, extensor hallucis, and extensor digitorum muscle strengths were 1/5 with reduction of plantar flexion and stepping gait. Additionally, a cystic mass that was approximately 5 cm long and 3 cm width was palpable at the proximal neck of fibula. Medical record revealed that swelling had occurred 2 months before and pain with motor deficit progressed rapidly during days. There was no trauma history. He had been diagnosed with lumbar magnetic resonance imaging (MRI), however no pathologic findings had been found. Therefore, electromyography (EMG), MRI, and soft tissue ultrasound (US) were performed. A hypodense mass around the fibular neck extending distally to syndesmosis was detected with MRI (Fig. 1a and b). Decreased motor potential amplitude and a loss of conductivity in the region of the proximal fibular head were seen in EMG study. These EMG findings were relevant to acute denervation of deep peroneal nerve. US showed a cystic, hypoechogenic lesion over the proximal fibular head. Excisional biopsy was performed with provisional diagnosis of synovial cyst (Fig. 2). Surgical exploration revealed synovial cyst that applied pressure to deep peroneal nerve in the 1/3 proximal part of cruris. The presence of hyaluronic acid in the synovial cyst was detected in the histologic examination (Figs. 3 and 4). Motor functions had started to recover within the

* Corresponding author at: Bezmialem Universitesi, Vakıf Gureba Hastanesi Ortopedi ve Travmatoloji AD, 34093 Fatih, Istanbul, Turkey. Tel.: +90 5324249732; fax: +90 2124531700.

E-mail addresses: drmehmeterdil@gmail.com, mehmeterdil@hotmail.com (M. Erdil), korhanozkan@hotmail.com (K. Ozkan), feyzaunlu@hotmail.com (F.U. Ozkan), kbilsel@gmail.com (K. Bilsel), dr.ismailturkman@gmail.com (I. Turkmen), serkansenol@hotmail.com (S. Senol), serhansarar@hotmail.com (S. Sarar).




Fig. 1. Preoperative MRI views: (a) coronal view and (b) axial view.

first week of operation. At the 2nd month of surgery total recovery of sensorial and motor functions were seen. Re-innervation of deep peroneal nerve was detected in EMG study at the 2nd month follow-up.

3. Discussion

Focal nerve entrapment syndromes due to ganglia are not common entities especially in the lower limb. List of peroneal nerve palsies include prolonged squatting, infection, varicose veins, rapid marked weight reduction, schwannoma, nerve herniation through a facial defect, neurofibromatosis, pneumatic compression, knee arthroplasty, high tibial osteotomy, and synovial cysts, to name a few.^{5–15} We described a deep peroneal neuropathy due to compression of a synovial cyst.

Compression to peroneal nerve by a synovial cyst was first described in 1921 by Sultan.⁴ Additionally, several cases of intra-neural synovial cysts of peroneal nerve have been reported.^{16–18} Reports of deep peroneal nerve compression due to a synovial cyst are much less.^{5,19,20}

Synovial cysts are mucinous filled cysts that can be found adjacent to joint capsule or tendon sheath. They can originate within perineural tissue or in an adjacent joint space or bursa and extend toward the nerve.²¹ They consist of an outer fibrous coat and an inner synovial lining and contain a clear, colorless, gelatinous fluid.²¹ In our case we observed these findings during the surgery (Fig. 2). Additionally, histologic examination revealed presence of hyaluronic acid in the synovial cyst (Figs. 3 and 4).

The frequent non-specific symptoms and findings like pain, swelling cause difficulty in differential diagnosis of synovial sarcoma, extra-skeletal chondrosarcoma, venous aneurysm. Diagnosis can be confirmed with imaging studies and ultimately surgical exploration.²² US is a non-invasive and cheap screening method, however, it is not sensitive enough to distinguish ganglia from other nerve sheath tumors. MRI can demonstrate fluid filled cystic




Fig. 3. Histopathologic view.




Fig. 4. Histopathologic view.

lesion that is hypointense on T1 and hyperintense on T2-weighted images.^{22,23} Electrophysiological tests can help to localize the lesion along the course of the nerve and can help to distinguish entrapment of deep peroneal nerve from sciatic neuropathy or L5 radiculopathy.^{24–26} In our case, we performed EMG, MRI and US to confirm the diagnosis preoperatively. MRI showed a hypodense mass around the fibular neck (Fig. 1), US showed a cystic, hypoechogenic lesion over the proximal fibular head, and EMG showed decreased motor potential amplitude and a loss of conductivity in the region of the proximal fibular head.

Synovial cysts that cause neuropathy can be successfully treated with surgical removal of the cyst. Recurrence rate is reported as approximately 10%. In order to prevent the recurrence, removal of the stalk and the base of the cyst in the tibiofibular joint could be performed.²⁷ We performed surgical excision of the cyst with removing the stalk and the base of the cyst in the tibiofibular joint. At 14th month follow-up, there were no findings of recurrence.




Fig. 2. Intraoperative views of the patient: (a) deep peroneal nerve beneath the synovial cyst, (b) ganglion stalk extending toward the tibiofibular joint, and (c) excised cyst.

This case demonstrates the etiologic cause of sudden drop foot and demonstrates the importance of considering synovial cyst around the knee in the differential diagnosis of deep peroneal neuropathy.

Conflicts of interest

None.

Funding

None.

Ethical approval

Obtained.

Author contributions

Mehmet Erdil, corresponding author of the case, diagnosed the patient and made study design. Korhan Ozkan had performed the surgery with Dr. Erdil, and also worked in data collecting (follow-up data). Feyza Unlu Ozkan, helped in English language editing and writing the manuscript. Kerem Bilisel, had attended the surgery with Dr. Erdil and Dr. Ozkan. He had taken the intraoperative photos. Ismail Turkmen worked in data collecting (follow-up data) with Dr. Ozkan. Serkan Senol worked in data collecting by preparing and evaluating the histopathologic specimens. Serhan Sarar had been in the surgery team as anesthesiologist, also he had helped in preparing the images of the manuscript.

References

- Erkin G, Uysal H, Keles I, Aybay C, Ozel S. Acute ulnar neuropathy at the wrist: a case report and review of the literature. *Rheumatology International* 2006;**27**:191–6.
- Strickland JW, Steichen JB. Nerve tumors of the hand and forearm. *Journal of Hand Surgery American Volume* 1977;**2**:285–91.
- Kerrigan JJ, Bertoni JM, Jaeger SH. Ganglion cysts and carpal tunnel syndrome. *Journal of Hand Surgery American Volume* 1988;**13**:763–5.
- Sultan C. Ganglion der Nervenscheide des Nervus Peroneus. *Zentralblatt für Chirurgie* 1921;**48**:963–5.
- Mont MA, Dellon AL, Chen F, Hungerford MW, Krackow KA, Hungerford DS. The operative treatment of peroneal nerve palsy. *Journal of Bone and Joint Surgery: American Volume* 1996;**78**:863–9.
- Yilmaz E, Karakurt L, Serin E, Güzel H. Peroneal nerve palsy due to rare reasons: a report of three cases. *Acta Orthopaedica et Traumatologica Turcica* 2004;**38**:75–8.
- Sangwan SS, Marya KM, Kundu ZS, Yadav V, Devgan A, Siwach RC. Compressive peroneal neuropathy during harvesting season in Indian farmers. *Tropical Doctor* 2004;**34**:244–6.
- Schulze J, Troeger H. Fibular nerve compression due to an unusual cause. A case report. *Handchirurgie, Mikrochirurgie, Plastische Chirurgie* 2004;**36**:25–8.
- Yamamoto N, Koyano K. Neurovascular compression of the common peroneal nerve by varicose veins. *European Journal of Vascular and Endovascular Surgery* 2004;**28**:335–8.
- Shahar E, Landau E, Genizi J. Adolescence peroneal neuropathy associated with rapid marked weight reduction: case report and literature review. *European Journal of Paediatric Neurology* 2007;**11**:50–4.
- Mahitchi E, Van Linthoudt D. Schwannoma of the deep peroneal nerve. An unusual presentation in rheumatology. *Praxis (Bern 1994)* 2007;**96**:69–72.
- Yang LJ, Gala VC, McGillicuddy JE. Superficial peroneal nerve syndrome: an unusual nerve entrapment. Case report. *Journal of Neurosurgery* 2006;**104**:820–3.
- Cebesoy O, Tutar E, Isik M, Arpaciglu O. A case of isolated giant plexiform neurofibroma involving all branches of the common peroneal nerve. *Archives of Orthopaedic and Trauma Surgery* 2007;**127**:709–12.
- Fukuda H. Bilateral peroneal nerve palsy caused by intermittent pneumatic compression. *Internal Medicine* 2006;**45**:93–4.
- Dallari D, Pellacani A, Marinelli A, Verni E, Giunti A. Deep peroneal nerve paresis in a runner caused by ganglion at capitulum peronei. Case report and review of the literature. *Journal of Sports Medicine and Physical Fitness* 2004;**44**:436–40.
- Spinner RJ, Ibrahim Elshiekh MA, Tubbs RS, Turner 3rd NS, Amrami KK. Posttraumatic torsional injury as an indirect cause of fibular intraneural ganglion cysts: case illustrations and potential mechanisms. *Clinical Anatomy* 2012;**25**(July (5)):641–6. <http://dx.doi.org/10.1002/ca.21290>.
- Al Mufargi YS, Mouch CA, Ziebarth K, Joeris A, Slongo T. An unusual cause of paralysis of the peroneal nerve: a report of 3 cases. *Journal of Pediatric Orthopedics* 2011;**31**(July–August (5)):e50–2.
- Waldschmidt U, Slongo T. An unusual cause of paralysis of the peroneal nerve—a case report. *Journal of Pediatric Surgery* 2010;**45**(January (1)):259–61.
- Rubin DI, Nottmeier E, Blasser KE, Peterson JJ, Kennelly K. Acute onset of deep peroneal neuropathy during a golf game resulting from a ganglion cyst. *Journal of Clinical Neuromuscular Disease* 2004;**6**(December (2)):49–53.
- Wadstein T. Two cases of ganglia in the sheath of the peroneal nerve. *Acta Orthopaedica Scandinavica* 1931;**2**:221–31.
- Mahaley MS. Ganglion of the posterior tibial nerve: case report. *Journal of Neurosurgery* 1974;**40**:120–4.
- Masciocchi C, Innacoli M, Cisternino S, Barile A, Rossi F, Passariello R. Myxoid intraneural cysts of external popliteal ischiadic nerve. Report of 2 cases studied with ultrasound, computed tomography and magnetic resonance imaging. *European Journal of Radiology* 1992;**14**:52–5.
- Leon J, Marano G. MRI of peroneal nerve entrapment due to a ganglion cyst. *Magnetic Resonance Imaging* 1987;**5**:307–9.
- Gutmann L. Atypical deep peroneal neuropathy in presence of accessory deep peroneal nerve. *Journal of Neurology, Neurosurgery and Psychiatry* 1970;**33**:453–6.
- Ciucci G, Callegarini C, Stumpo M, Poppi M. Ganglionic cyst of the deep peroneal nerve: description of a case. *Italian Journal of Neurological Sciences* 1996;**17**:83–6.
- Antonini G, Bastianello S, Nucci F, Artico M, Bozzao L, Millefiorini M. Ganglion of deep peroneal nerve: electrophysiology and CT scan in the diagnosis. *Electromyography and Clinical Neurophysiology* 1991;**31**:9–13.
- Muckart RD. Compression of the common peroneal nerve by intramuscular ganglion from the superior tibio-fibular joint. *Journal of Bone and Joint Surgery: British Volume* 1976;**58**:241–4.