



Scleredema of Buschke in a little child and its successful treatment with ultraviolet A1 phototherapy

Küçük bir çocukta Buschke'nin sklerödemi ve ultraviyole A1 fototerapisi ile başarılı tedavisi

• Damla Demir, • Müge Göre Karaali*, • Filiz Cebeci Kahraman**, • İlkin Zindancı***, • Ebru Zemheri****

Istinye University Faculty of Medicine, Department of Dermatology, İstanbul, Turkey

*İrmet International Hospital, Clinic of Dermatology, Tekirdağ, Turkey

**Medeniyet University Faculty of Medicine, Department of Dermatology, İstanbul, Turkey

***University of Health Sciences Turkey, Hamidiye Faculty of Medicine, Ümraniye Training and Research Hospital, Clinic of Dermatology;

****Clinic of Pathology, İstanbul, Turkey

Abstract

A 5-year-old girl with skin hardness and thickening was admitted to our clinic and diagnosed with scleredema of Buschke clinically and histopathologically. Ultraviolet A1 (UVA1) phototherapy at a dose of 30 J/cm² was planned for 3 days in 1 week. After 45 UVA1 phototherapy sessions (with a cumulative dose of 1350 J/cm²), clinical complete remission was achieved. Ultrasonography revealed improvement in skin thickening, and no recurrence was noted in the 2-year follow-up period.

Keywords: Scleredema of Buschke, UVA1, phototherapy, scleredema, scleredema adultorum, ultrasonography

Öz

Beş yaşında kadın hasta, deride sertlik ve kalınlık şikayeti ile kliniğimize başvurdu. Klinik ve histopatolojik bulgular ile Buschke'nin sklerödemi tanısı kondu. Haftada 3 gün 30 j/cm² ultraviyole A1 (UVA1) fototerapisi planlandı. Kırk beş seans UVA1 fototerapisinden sonra (kümülatif doz 1,350 j/cm² ile) klinik tam remisyona sağlandı. Ultrasonografi ile deri kalınlığında iyileşme gözlemlendi ve 2 yıllık takipte nüks izlenmedi.

Anahtar Kelimeler: Buschke'nin sklerödemi, UVA1, fototerapi, skleredema, sklerema adultorum, ultrasonografi

Introduction

Scleredema of Buschke (SB) is a rare connective tissue disease and is characterized by skin induration. It typically starts from the neck and spreads to the upper half of the trunk, shoulders, back, and face¹. It was first described by Curzio in 1752, but was named after Abraham Buschke, who reported a 46-year-old male patient who presented with skin

thickening and hardening after an influenza epidemic in 1902².

SB has three types. Type 1 occurs following infectious diseases and tends to improve spontaneously. Type 2 is more persistent and associated with paraproteinemia, not with infectious diseases. Type 3 occurs in patients with diabetes mellitus and has a male predominance, unlike other types, and it is also called scleredema diabetorum³.

Address for Correspondence/Yazışma Adresi: Müge Göre Karaali MD, İrmet International Hospital, Clinic of Dermatology, Tekirdağ, Turkey

Phone: +90 530 309 33 28 **E-mail:** mugegore@hotmail.com **Received/Geliş Tarihi:** 24.06.2022 **Accepted/Kabul Tarihi:** 31.10.2022

ORCID: orcid.org/0000-0003-0645-0662

The study was conducted in Department of Dermatology, University of Health Sciences Turkey, Hamidiye Faculty of Medicine, Ümraniye Training and Research Hospital, İstanbul, Turkey.

Cite this article as: Demir D, Göre Karaali M, Cebeci Kahraman F, Zindancı İ, Zemheri E. Scleredema of Buschke in a little child and its successful treatment with ultraviolet A1 phototherapy. Turkderm-Turk Arch Dermatol Venereol 2022;56:186-9

©Copyright 2022 by Turkish Society of Dermatology and Venereology
Turkderm-Turkish Archives of Dermatology and Venereology published by Galenos Yayınevi.



Patients with SB have abnormal extracellular matrix gene expression of fibroblasts, and the accumulation of collagen and aminoglycans in the dermis is the main disease mechanism⁴. Immunosuppressive agents, antibiotics, corticosteroids, radiotherapy, and phototherapy are treatment options, but with unsuccessful results¹. Possible treatment options are limited, especially in pediatric patients owing to their negative effects and side effects on growth and development. Herein, we present a child with SB treated with ultraviolet A1 (UVA1) phototherapy. In this report, we aimed to emphasize the effectiveness of UVA1 phototherapy, which is a safe and successful treatment method in the treatment of SB, especially in children.

Case Report

A 5-year-old girl presented to our outpatient clinic because of skin hardness and thickening for 4 months. She had no history of any additional disease. A dermatological examination revealed woody hardness and skin thickening predominantly on her face and upper and lower extremities (Figure 1a-c). Laboratory tests were not remarkable. A 4 mm punch biopsy was performed, and histopathological examination revealed loss of hair follicles in the dermis, enlarged collagen bundles (Figure 2a, b), and mucin deposition between collagen fibers (Figure 2c). She was diagnosed with SB based on the clinical and histopathological findings.

Before treatment, skin thickness was evaluated by ultrasonography (Figure 3a, c). UVA1 phototherapy was planned for 3 days in 1 week. A dose of 30 J/cm² UVA1 was applied for each treatment session. After 45 sessions of UVA1 phototherapy (with a cumulative dose of

1350 J/cm²), clinical complete remission was achieved, and treatment was concluded. Ultrasonography revealed improvement of the skin thickening (Figure 3b, d). No recurrence was noted in the 2-year follow-up period.

The UVA1 irradiation equipment consisted of a Waldmann 7001 K cabin with Waldmann TL10R low-pressure lamps (Waldmann GmbH, Schwenningen, Germany). These lamps generate UVA1 wavelengths in the 340-400 nm range. Infrared irradiation is also emitted, but this is filtered by an acrylic glass screen. The UVA1 irradiation level is approximately 35 mW/cm². A dose of 30 J/cm² is achieved in approximately 30 min.

Discussion

UVA1 is a part of the UV spectrum that emits photons with wavelengths of 340-400 nm⁵. The two main sources of UVA1 are fluorescent lamps and metal halides. Fluorescent lamp cells only allow low dose (LD) (10-30 J/cm²) or medium dose (MD) (40-70 J/cm²). High-output, high-dose metal halide sources allow up to (130 J/cm²)⁶. UVA1 phototherapy was reported to be effective in the treatment of inflammatory skin diseases such as atopic dermatitis⁷. Mempel et al.⁸ observed that mature collagens 1 and 3 decreased in skin biopsies after UVA1 treatment in 14 patients with atopic dermatitis. In addition, in 1981, Mutzhas et al.⁵ showed that UVA wavelengths reached the deep dermis and subcutis. In later studies, UVA1 treatment was thought to be an effective treatment option for connective tissue diseases with sclerosis.

In vitro studies in cultured fibroblasts applying UVA1 phototherapy have shown an increase in collagenase production^{9,10}. However, UVA1



Figure 1. (a-c) Woody hardness and thickness of the skin predominantly on extremities




Figure 2. (a, b) Loss of hair follicles in the dermis, enlarged collagen bundles (hematoxylin and eosin staining, x10; Verhoeff Von Gieson staining, x10). **(c)** Mucin deposition between collagen fibers (periodic acid Schiff staining, x40)




Figure 3. (a) Pretreatment ultrasonographic skin thickness of the right arm: 1.5 mm, (b) post-treatment ultrasonographic skin thickness of the right arm: 1.2 mm, (c) pretreatment ultrasonographic skin thickness of the left arm: 1.5 mm, and (d) post-treatment ultrasonographic skin thickness of the left arm: 1.3 mm

reduces the enzyme activity of propyl hydroxylase, which stabilizes the triple helix structure of collagen, and causes disruption of the cross-linking between collagen fibrils^{11,12}. In addition, many *in vivo* and *in vitro* studies have shown that UVA increases the number of matrix metalloproteinase (MMP)-specific mRNA receptors in cultured fibroblasts. UVA1 stimulated three members of the MMP family, i.e., MMP1, MMP3, and MMP9, which are responsible for the degradation of type 1 collagen¹³. On the contrary, UVA1 reduces transforming growth factor (TGF)-beta cell surface receptors, which play an important role in procollagen synthesis, growth, and differentiation¹⁴. Recently, some studies have focused on SMAD, which is a family of transcription factor proteins involved in TGF-beta signaling from the cell surface to the nucleus. *In vitro* and *in vivo* studies have shown that SMAD7 is an important inhibitor of the TGF-beta SMAD pathway and UVA1 stimulates SMAD7^{15,16}. Based on these mechanisms of action, UVA1 phototherapy was reported as an effective treatment in different connective tissue diseases with sclerosis.

SB is a chronic skin disease with sclerosis. The effectiveness of UVA1 for the treatment of SB was first reported in 2004¹⁷. Two patients diagnosed with type 2 and type 3 SB were treated with LD UVA1 phototherapy, and regression was reported in all lesions. In 2005, Eberlein-König et al.¹⁸ reported thinning of the skin following MD UVA1 treatment in a patient diagnosed with type 3 SB. In a study retrospectively evaluating 92 patients with different skin diseases treated with UVA1 in 2006, five had SB, clinical regression was detected in four patients, and treatment was terminated in one patient because of polymorphous light eruption¹⁹. In our case, pre- and post-treatment ultrasonography findings were also evaluated. After 45 sessions of UVA1 treatment,

ultrasonography detected regression of the lesions and significant skin thinning. No treatment-related side effects occurred.

UVA1 phototherapy is an effective and safe treatment method for patients with SB. It may be a treatment option, especially in pediatric patients.

Ethics

Informed Consent: Retrospective study.

Peer-review: Externally and internally peer-reviewed.

Authorship Contributions

Surgical and Medical Practices: D.D., M.G.K., F.C.K., İ.Z., E.Z., Concept: D.D., M.G.K., F.C.K., İ.Z., E.Z., Design: D.D., M.G.K., F.C.K., İ.Z., E.Z., Data Collection or Processing: D.D., M.G.K., F.C.K., İ.Z., E.Z., Analysis or Interpretation: D.D., M.G.K., F.C.K., İ.Z., E.Z., Literature Search: D.D., M.G.K., F.C.K., İ.Z., E.Z., Writing: D.D., M.G.K., F.C.K., İ.Z., E.Z.

Conflict of Interest: No conflict of interest was declared by the authors.

Financial Disclosure: The authors declared that this study received no financial support.

References

1. Beers WH, Ince A, Moore TL: Scleredema adultorum of Buschke: a case report and review of the literature. *Semin Arthritis Rheum* 2006;35:355-9.
2. Busschke A: Vorstellung eines Falles von Sklerödem vor der Berliner Gesellschaft für Dermatologie. *Arch Dermatol Syph* 1900;25:283.
3. Scleredema adultorum. *Arch Dermatol* 1968;98:319-20.
4. Varga J, Gotta S, Li L, Di Leonardo M: Scleredema adultorum: case report and demonstration of abnormal expression of extracellular matrix genes in skin fibroblasts *in vivo* and *in vitro*. *Br J Dermatol* 1995;132:992-9.

5. Mutzhas MF, Holzle E, Hofmann C, Plewig G: A new apparatus with high radiation energy between 320-460 nm: physical description and dermatological applications. *J Invest Dermatol* 1981;76:42-7.
6. Dawe RS: Ultraviolet A1 phototherapy. *Br J Dermatol* 2003;148:626-37.
7. Meffert H: Phototherapy of skin diseases. *Z Arztl Fortbild* 1992;86:947-50.
8. Mempel M, Schmidt T, Boeck K, et al.: Changes in collagen I and collagen III metabolism in patients with generalized atopic eczema undergoing medium-dose ultraviolet A1 phototherapy. *Br J Dermatol* 2000;142:473-80.
9. Gruss C, Reed JA, Altmeyer P, McNutt NS, Kerscher M: Induction of interstitial collagenase (MMP-1) by UVA-1 phototherapy in morphed fibroblasts. *Lancet* 1997;350:1295-6.
10. Scharffetter K, Wlaschek M, Hogg A, et al.: UVA irradiation induces collagenase in human dermal fibroblasts in vitro and in vivo. *Arch Dermatol Res* 1991;283:506-11.
11. Johnston KJ, Oikarinen AJ, Lowe NJ, Clark JG, Uitto J: Ultraviolet radiation-induced connective tissue changes in the skin of hairless mice. *J Invest Dermatol* 1984;82:587-90.
12. Oikarinen A, Karvonen J, Uitto J, Hannuksela M: Connective tissue alterations in skin exposed to natural and therapeutic UV-radiation. *Photodermatol* 1985;2:15-26.
13. Fisher GJ, Kang S: Phototherapy for scleroderma: biologic rationale, results, and promise. *Curr Opin Rheumatol* 2002;14:723-6.
14. El-Mofty M, Mostafa W, Esmat S, et al.: Suggested mechanisms of action of UVA phototherapy in morphea: a molecular study. *Photodermatol Photoimmunol Photomed* 2004;16:43-9.
15. Dong C, Zhu S, Wang T, et al.: Deficient Smad7 expression: a putative molecular defect in scleroderma. *Proc Natl Acad Sci U S A* 2002;99:3908-13.
16. Moustakas A, Souchelnytskyi S, Heldin CH: Smad regulation in TGF-beta signal transduction. *J Cell Sci* 2001;114:4359-69.
17. Janiga JJ, Ward DH, Lim HW: UVA-1 as a treatment for scleroderma. *Photodermatol Photoimmunol Photomed* 2004;20:201-1.
18. Eberlein-König B, Vogel M, Katzer K, et al.: Successful UVA1 phototherapy in a patient with scleredema adultorum. *J Eur Acad Dermatol Venerol* 2005;19:203-4.
19. Tuchinda C, Kerr HA, Taylor CR, et al.: UVA1 phototherapy for cutaneous diseases: an experience of 92 cases in the United States. *Photodermatol Photoimmunol Photomed* 2006;22:247-53.