

MORPHOMETRIC AND GENETIC CHARACTERIZATION OF HONEY BEES (*APIS MELLIFERA* L.) FROM THRACE REGION OF TURKEY

Fulya Özdil^{1*}Devrim Oskay²Raziye Işık²Selen Yatkın²Abdurrahman Aydın³Ahmet Güler³¹Bioengineering, Istanbul Medeniyet University / Agricultural Biotechnology, Tekirdağ Namık Kemal University, Turkey²Agricultural Biotechnology, Tekirdağ Namık Kemal University, Turkey³Animal Science, Ondokuzmayıs University, Turkey

*corresponding author: fulya.ozdil@medeniyet.edu.tr, fozdil@nku.edu.tr

Received: 22 September 2021; accepted: 24 March 2022

Abstract

A detailed morphological and genetic characterization of honey bees from the Thrace and west Anatolian regions of Turkey was surveyed. A total of 1650 worker bee samples (110 colonies) were evaluated with the forty-one morphological characters and 217 honey bee samples were analyzed via DNA sequencing of the tRNA^{leu}-cox2 region. In this study, three different populations, Thrace (Tekirdağ, Kırklareli and Edirne provinces), Island Gökçeada, and western Anatolia were formed based on morphometrics, since the Marmara Sea has taken a very strong barrier role in this formation. The morphological similarity of the Thrace population was supported by the genetic analysis. The sequencing of the tRNA^{leu}-cox2 region revealed twenty-two different haplotypes, sixteen of which are novel. The C2d, macedonica-like haplotype, was the most widely found haplotype (48%) all around the Thrace region. Along with the C2d haplotype, previously published C2s, C2v, C2i, C2j, and C2h haplotypes, and the newly found haplotypes were also observed but less frequently. In this study, Thrace honey bees were found to more similar to *A. m. macedonica* through the mtDNA sequence analysis, whereas carnica-like honey bees were only found near the Istranca mountain ridges, Kırklareli province and macedonica-like honey bees all around the Thrace region. According to our results, some of the Thrace honey bee populations may be both *A. m. carnica* and *A. m. macedonica* but the assignment to the latter subspecies seems more likely due to its geographic range.

Keywords: *Apis mellifera*, DNA sequencing, morphometrics, thrace honey bees, tRNA^{leu}-Cox2 gene

INTRODUCTION

The western honey bee, *Apis mellifera* L., has a wide distribution in Africa, the Middle East and Europe. Since Ruttner's morphological classification of twenty-five taxonomic groups (subspecies) of the western honey bee (*Apis mellifera* L.) (Ruttner, 1988), twenty-nine recognized subspecies of *Apis mellifera* has been identified with the addition of newly described subspecies including *A. m. ruttneri* from Malta (Sheppard et al., 1997), *A. m. pomonella* from

Central Asia (Sheppard & Meixner, 2003), *A. m. simensis* from Ethiopia (Meixner et al., 2011) and *Apis mellifera sinisxinyuan* Xinyuan bees from China (Chen et al., 2016). These subspecies are now typically divided into four major groups, supported by morphometric and molecular studies: A lineage in Africa, M lineage in western and northern Europe, C lineage in eastern Europe and O lineage in the Middle East and Western Asia (Garnery et al., 1992; Whitfield et al., 2006; Oleksa et al., 2011; Han et al., 2012; Péntek-Zakar et al., 2015). The existence of a

fifth lineage (Y) in the north-east side of Africa from the country of Ethiopia has also been proposed (Franck et al., 2001).

Turkey is in the contact zone of the different lineages and was first proposed to be O lineage, but then studies revealed it to be mostly C-lineage (Smith et al., 1997; Palmer et al., 2000; Kandemir et al., 2006; Özgül et al., 2009). Within Turkey, wide ranges of climates and habitats are found, and several honey bee subspecies and ecotypes have been described. According to Ruttner (1988), *A. m. anatoliaca*, *A. m. caucasica*, and *A. m. meda* exist in Turkey. Nearly all of the Turkey is occupied by the Anatolian honey bee (*A. m. anatoliaca*). *A. m. caucasica* and *A. m. meda* are found in the northeast and the southeast, respectively. Recent molecular studies of Turkish honeybees have shown that *A. m. syriaca* is found in the southern part of the country near the Hatay province (Palmer et al., 2000; Kandemir et al., 2006; Bodur et al., 2007; Solorzano et al., 2009) and in contrast to Ruttner's studies, morphometric (Güler & Kaftanoğlu, 1999; Güler, 2010; Güler et al., 2010) and molecular studies (mtDNA and microsatellites) have revealed a fifth honey bee subspecies in the Thrace region of Turkey (Smith et al., 1997; Palmer et al., 2000; Kandemir et al., 2006; Bodur et al., 2007; Kekecoglu et al., 2009; Ünal & Özgül, 2018). Thrace honey bee populations are considered as a different ecotype of *A. m. carnica* in Turkey's honey bee gene resources and registered officially by the Republic of Turkey, Ministry of Agriculture and Forestry (Anonymous, 2020).

Within the Balkan Peninsula very near to the Thrace region, several honey bee subspecies and ecotypes including *A. m. carnica*, *A. m. macedonica* and *A. m. cecropia* have been described through morphometrics and molecular studies but *A. m. carnica* is the most widely found pioneer honey bee subspecies in this region and the Carniolan subspecies was reported to have the most strains in the world in its natural geography (Sušnik et al., 2004; Bouga et al., 2005, 2011; Nedić et al., 2009; Muñoz et al., 2009, 2012; Martimianakis et al., 2011; Coroian et al., 2014; Meixner et al., 2014; Tanasković et al., 2021).

In this study, a detailed morphological and genetic characterization of honey bees from the Thrace and western Anatolian region of Turkey was surveyed. We made a comprehensive sampling and used forty-one morphological characteristics and sequenced the tRNA^{leu}-Cox2 intergenic region to identify the honey bee (*A. mellifera*) populations in the European part of Turkey-Thrace region (Tekirdağ, Kırklareli and Edirne provinces), Island Gökçeada and Çanakkale province (both Gallipoli Peninsula and western Anatolian side).

The aim of this study was threefold: (i) to provide a detailed morphological characterization of the Thrace and the western Anatolian region of Turkey, with a large sampling size through the analysis of morphological characteristics; (ii) to reveal the different haplotypes after the completion of the survey of mtDNA variation of the tRNA^{leu}-cox2 region through the sequencing of a large collection of individuals sampled across the entire Thrace and western Anatolian region and to revise the C lineage haplotypes which have already been deposited to the GenBank database; (iii) to determine whether the dominant Thrace region ecotype belonged to *A. m. caucasica*, *A. m. macedonica* or *A. m. carnica* subspecies.

MATERIAL AND METHODS

Sampling of the honey bees

The worker bees were collected between May 2014 and September 2015 for morphological and mitochondrial surveys. Honey bees were sampled from five different localities; three provinces in the Thrace region: Tekirdağ, Kırklareli, Edirne provinces; the Çanakkale province: (Gallipoli Peninsula and western Anatolian side); the island of Gökçeada on the Marmara Sea near Gallipoli Peninsula (Fig. 1). A total of 1650 worker bee samples (110 colonies from twenty-three apiaries, fifteen repetitions per colony) were evaluated for forty-one morphological characteristics. On the other hand, 217 worker bees, including morphologically measured 110 samples, from 133 apiaries in seventy-nine different localities with the two



Fig. 1. Sampling locations of honey bees in European part of Turkey-Thrace region (Tekirdağ, Kırklareli and Edirne), Island of Gökçeada and Çanakkale province (both Gallipoli Peninsula and western Anatolian side).

of each reference samples (*A. m. caucasica*, *A. m. carnica*, *A. m. macedonica*) were analysed for genetic characterization. One sample was taken from each colony, which derived from established colonies maintained by local non-migratory beekeepers. These samples were used for DNA sequencing of the tRNA^{eu}-cox2 region in the mitochondrial genome.

Morphological evaluation

The worker bee samples were collected and stored in 72% ethyl alcohol until preparation. Each part of the samples was fixed on the slide with chloral hydrate-free Hoyer's liquid. Each of the 110 colonies consisted of fifteen worker bees, so a total of 1650 (15x110) worker bees were used for morphological measuring. The right forewings of the fifteen worker bees from each colony were mounted on glass slides with Hoyer's liquid. In each sample, the standard and most commonly used anterior wing vein angles A_4 , B_4 , D_7 , E_9 , G_{18} , J_{10} , J_{16} , K_{19} , L_{13} , N_{23} and O_{26} were measured biometrically with the use of the software available in the Olympus trinocular stereo microscope (SZ61) (Ruttner et al., 1978; Moritz, 1991). Forty-one morphological characters given in Tab. 1 and Tab. 2 were measured as described by Moritz (1991), Güler

(2010) and Güler et al. (2010). Measurements were made with the Olympus SZ61 trinocular stereo microscope with the computer program (cellSens standart 1.8).

Statistical analysis

The morphometrics data were analysed through ANOVA, and the differences between the means were compared with the Student Newman Keuls (SNK) post hoc tests (SPSS, 2004). In addition, the Multivariate Discriminant Stepwise Analysis Method (MDSAM), which determines the differences and the grouping levels in terms of the morphological characteristics between more than two biological sources, was used to determine the Standard Multivariate Canonical Discriminant Function and Constant Descriptive Coefficients (SMCDFCDC) of the five different regions from Thrace and western Anatolia (Cooley & Lohnes, 1971). The territorial regions of the groups in a Coordinate system were determined and standardized with the use of the SMCDFCDC (Fig. 2).

Genetic Characterization

DNA isolation

The worker bees were individually placed in the Eppendorf tubes containing 95% ethanol

and transported to the Molecular Genetics Laboratory, the Department of Agricultural Biotechnology, Tekirdağ Namık Kemal University. Total genomic DNA was extracted from thoraces according to the phenol-chloroform extraction method (Sambrook & Russel, 2001). The genomic DNA concentration was quantified with a Qubit™ 2.0 Fluorometer (ThermoFisher Scientific), and 20-30 ng of genomic DNA was used for the PCR.

DNA sequencing

The tRNA^{leu}-cox2 gene region was amplified according to Garnery et al. (1992) with E2 and H2 primers. The amplified PCR products were sequenced on an Applied Biosystems 3500XL Genetic Analyzer (Applied Biosystems, USA) in order to verify the nucleotide variations. The sequences were assembled with ChromasPro version 1.7.6 and aligned with the BioEdit Sequence Alignment Editor with Clustal W multiple alignment modules (Hall, 1999). Phy-

Table 1.

The mean and standard errors of the directly measured morphological characters of the Thrace, Çanakkale and island Gökçeada worker bee samples

Character	Regions					X±Sx
	Tekirdağ	Kırklareli	Edirne	Çanakkale	Gökçeada	
LH***	0.238±0.004 ^c	0.228± 0.003 ^c	0.231±0.002 ^c	0.270±0.003 ^b	0.283±0.004 ^a	0.247±0.003
WTa***	1.117±0.022 ^b	1.100±0.015 ^{bc}	1.051±0.010 ^c	1.127±0.227 ^b	1.186±0.015 ^a	1.109±0.009
WTb*	0.300±0.012 ^{ab}	0.316±0.011 ^{ab}	0.305±0.011 ^{ab}	0.336±0.012 ^a	0.296±0.013 ^b	0.311±0.006
LPr**	6.247±0.077 ^b	6.316±0.072 ^{ab}	6.319±0.023 ^{ab}	6.511±0.068 ^a	6.422±0.108 ^{ab}	6.351±0.030
LF ^{NS}	2.621±0.016	2.637±0.011	2.622±0.008	2.631±0.020	2.663±0.016	2.633±0.006
LT*	3.144±0.019 ^{ab}	3.145±0.016 ^{ab}	3.116±0.011 ^b	3.145±0.023 ^{ab}	3.178±0.016 ^a	3.143±0.008
LM*	2.044±0.013 ^b	2.092±0.013 ^a	2.078±0.009 ^{ab}	2.073±0.013 ^{ab}	2.092±0.013 ^a	2.074±0.006
WM*	1.168±0.008 ^{ab}	1.162±0.009 ^b	1.190±0.006 ^a	1.169±0.009 ^{ab}	1.155±0.008 ^b	1.170±0.004
WT ₃ ***	1.896±0.007 ^b	1.869±0.008 ^b	1.877±0.007 ^b	1.961±0.013 ^a	1.984±0.015 ^a	1.910±0.006
WT ₄ ***	1.846±0.007 ^c	1.806±0.008 ^d	1.822±0.007 ^{cd}	1.920±0.012 ^b	1.953±0.014 ^a	1.859±0.006
WS ₃ *	2.811±0.012 ^b	2.804±0.014 ^b	2.792±0.009 ^b	2.912±0.076 ^a	2.823±0.013 ^{ab}	2.825±0.015
LWM***	1.481±0.020 ^b	1.421±0.009 ^{bc}	1.404±0.006 ^c	1.664±0.043 ^a	1.639±0.009 ^a	1.505±0.014
WWM ^{NS}	2.346±0.012	2.361±0.013	2.342±0.011	2.396±0.057	2.342±0.013	2.357±0.012
DWM***	0.315±0.005 ^b	0.325±0.005 ^b	0.317±0.005 ^b	0.349±0.011 ^a	0.352±0.006 ^a	0.329±0.003
LS ₆ U*	2.606±0.012 ^b	2.633±0.014 ^{ab}	2.640±0.010 ^{ab}	2.642±0.031 ^{ab}	2.679±0.017 ^a	2.637±0.008
WS ₆ *	3.119±0.018 ^b	3.137±0.018 ^b	3.128±0.015 ^b	3.161±0.041 ^{ab}	3.215±0.016 ^a	3.146±0.011
LFW***	8.333±0.022 ^b	8.343±0.036 ^b	8.362±0.018 ^b	8.546±0.016 ^a	8.486±0.028 ^a	8.402±0.014
WFW**	2.856±0.009 ^b	2.866±0.015 ^{ab}	2.878±0.008 ^{ab}	2.902±0.008 ^{ab}	2.893±0.019 ^a	2.877±0.006
LCa**	0.491±0.004 ^{ab}	0.504±0.005 ^a	0.502±0.004 ^a	0.488±0.005 ^{ab}	0.484±0.005 ^b	0.495±0.002
LCb***	0.218±0.003 ^{ab}	0.212±0.006 ^{bc}	0.201±0.002 ^c	0.230±0.003 ^a	0.226±0.003 ^a	0.216±0.002
CT ₂ ***	6.981±0.105 ^a	6.788±0.105 ^{ab}	6.597±0.103 ^b	5.994±0.153 ^c	6.615±0.125 ^b	6.621±0.060
CT ₃ **	6.763±0.111 ^a	6.403±0.092 ^b	6.389±0.078 ^b	6.099±0.207 ^b	6.787±0.108 ^a	6.479±0.059
CT ₄ ***	4.200±0.069 ^a	4.365±0.056 ^a	4.372±0.064 ^a	3.580±0.103 ^c	3.831±0.108 ^b	4.113±0.045
CSc**	0.853±0.147 ^a	0.451±0.078 ^{bc}	0.325±0.059 ^c	0.863±0.109 ^a	0.666±0.135 ^{ab}	0.618±0.052

Length of hairs (LH), Width tomentum a (WTa), Width tomentum b (WTb), Length of proboscis (LPr), Length of femur (LF), Length of tibia (LT), Length of metatarsus (LM), Width of metatarsus (WM), , Width of tergite 3 (WT3), Width of tergite 4 (WT4), Width of sternit 3 (WS3), Length of wax mirror, (LWM), Width of wax mirror (WWM), D. Between mirrors (DWM), Length of sternum 6 (LS6), Width of sternum 6 (WS6), Length of forewing (LFW), Width of forewing (WFW), Length of cubital a (LCa), Length of cubital b (LCb), Colour of tergite 2 (CT₂), colour of tergite 3 (CT₃), Colour of tergite 4 (CT₄), Colour of scutellum (CSc)

NS, *, **, *** non significant and significant at 0.05, 0.01, and 0.001 respectively.

logenetic trees were generated with the use of the neighbor-joining algorithm in MEGA 6 (Tamura et al., 2013). The SplitsTree was also used to construct a network from the distance matrices based on allele-sharing distances (Huson & Bryant, 2006).

RESULTS

The worker-bee samples belonging to the five locations were found to be similar in only five characters but different from one another ($P < 0.001$) in thirty-six characters (Tab. 1 and Tab. 2).

The Discriminant Analysis Stepwise method showed that the hair length (HL), wing L_{13} vein angle, fourth tergite width (T_4), wing N_{23} vein angle, metatarsal index (MI), wing E_9 vein angle, second tergite color (CT_2), scutellum color (CS_C), both wing O_{26} and D_7 vein angles and wing length (WL) characters determined that the worker-bee samples represented the different ($P < 0.001$) bee populations respectively (Tab. 1 and Tab. 2).

The method determined 94.5% to be the proper grouping level of 110 worker bee samples representing these five different locations. Grouping in different areas indicated that these bees come from different genetic sources in terms of morphological structure Tab. 3.

Morphological differences according to the discriminant functions (F1) were found among the worker bee samples of the Thrace Region (Kırklareli, Tekirdağ and Edirne), Çanakkale and Gökçeada Island (Fig. 2). The Çanakkale and Gökçeada samples formed a cluster in completely different areas from one another and from samples of the Thrace Region (100%). Therefore, three different populations were formed in terms of the morphological structure: one on the continent of the Europe-Thrace region, one in the Çanakkale province on Asian/Anatolian side of the Turkey and one Gökçeada Island. The Sea of Marmara and Gallipoli Peninsula, important barriers in separating Asian and European continents, have also played an important role in forming this genetic variation.




Fig. 2. Discriminant Analysis of honey bee (*Apis mellifera*) samples from different genotypes (Tekirdağ, Edirne, Kırklareli, Çanakkale and Gökçeada Island). Horizontal axis: canonical function 1; vertical axis: canonical function 2.

Table 2.

The mean and standard errors of some calculated and direct measured morphological characters of the Thrace, Çanakkale and Gökçeada regions worker bee samples

Character	Regions				X±Sx
	Tekirdağ	Kırklareli	Edirne	Gökçeada	
TI*	4.284±0.232 ^{ab}	3.887±0.148 ^b	3.809±0.165 ^b	4.683±0.231 ^a	4.080±0.093
LHL ^{NS}	7.809±0.045	7.871±0.033	7.816±0.027	7.934±0.043	7.846±0.018
CI***	2.302±0.043 ^b	2.471±0.044 ^a	2.552±0.051 ^a	2.188±0.048 ^b	2.365±0.025
T ₃ +T ₄ ***	3.737±0.015 ^c	3.675±0.017 ^d	3.695±0.014 ^{cd}	3.937±0.026 ^a	3.765±0.013
MI***	57.206±0.387 ^a	55.716±0.331 ^{bc}	57.394±0.317 ^a	55.236±0.194 ^c	56.508±0.180
S ₆ ^{NS}	83.398±0.490	84.056±0.296	84.581±0.327	83.452±0.301	83.875±0.182
A ₄ ***	32.109±0.206 ^{bc}	30.970±0.242 ^d	31.769±0.249 ^c	32.532±0.193 ^{ab}	31.953±0.122
B ₄ ***	104.037±0.543 ^{ab}	105.727±0.598 ^a	103.399±0.660 ^{bc}	101.882±0.517 ^{cd}	103.474±0.312
D ₇ ***	100.857±0.344 ^a	99.495±0.377 ^b	99.413±0.355 ^b	101.121±0.376 ^a	100.432±0.193
E ₉ ***	20.392±0.153 ^{bc}	21.013±0.153 ^a	20.693±0.252 ^{ab}	19.158±0.294 ^d	20.336±0.105
G ₁₂ **	88.837±0.430 ^{bc}	88.360±0.348 ^{bc}	87.933±0.348 ^c	89.480±0.436 ^{ab}	88.857±0.184
J ₁₀ **	52.605±0.398 ^b	53.989±0.358 ^a	54.274±0.443 ^a	52.536±0.520 ^b	53.521±0.206
J ₁₆ ***	90.103±0.385 ^{bc}	91.953±0.357 ^a	90.859±0.401 ^{ab}	89.436±0.434 ^c	90.936±0.196
K ₁₉ ^{NS}	74.372±0.367	75.440±0.416	75.356±0.291	74.662±0.499	75.104±0.179
L ₁₃ ***	16.145±0.119 ^b	15.556±0.143 ^{cd}	15.390±0.112 ^d	15.989±0.273 ^{bc}	16.112±0.112
N ₂₃ ***	88.683±0.402 ^{cd}	91.161±0.263 ^b	89.854±0.405 ^{bc}	88.180±0.408 ^d	90.212±0.288
O ₂₆ *	34.496±0.420 ^b	35.195±0.456 ^{ab}	34.147±0.444 ^b	36.538±0.807 ^a	34.862±0.239

Tomentum Index (TI), Length of Hind Leg (LHL), Cubital Index (CI), Body size (T₃+T₄), Metatarsal Index (MI), Sternum 6 Index (S₆), Veinal angles A₄, B₄, D₇, E₉, G₁₂, J₁₀, J₁₆, K₁₉, L₁₃, N₂₃, O₂₆

NS, *, **, *** non significant and significant at 0.05, 0.01, and 0.001 respectively.

The populations of the Thrace region (Tekirdağ, Edirne and Kırklareli) were found similar to one another in relation to the morphological structure. The samples of the region were overlapped at certain levels. For example, two samples (8%) from Tekirdağ were placed within

Table 3.

Grouping levels of 110 worker bee samples collected from bee genotypes in five different areas and the number and ratios of the genotype groups each sample represents

	Groups Tekirdağ Kırklareli	Estimation group memberships			Total
		Edirne	Çanakkale	Gökçeada	
Oriijinal Groups	Tekirdağ	23	2		25
	Kırklareli		23	2	25
	Number Edirne		2	23	25
	Çanakkale			20	20
	Gökçeada			15	15
	Tekirdağ	92.0	8.0		100.0
	Kırklareli		92.0	8.0	100.0
	% Edirne		8.0	92.0	100.0
	Çanakkale			100.0	100.0
	Gökçeada			100.0	100.0

the samples from Kırklareli and two samples (8%) from Kırklareli in Edirne, and two samples (8%) from Edirne in Kırklareli. The overlapped clustering (Fig. 2) in this region was the result of a similar morphological structure. In fact, the worker bee samples in the Thrace region were found to be similar to one another in some of the morphological characteristics (HL, WT₃, WT_b, WS₃, DWM, LS₆, WS₆, LFW, and CT₄) (Tab. 1 and Tab. 2). On the other hand, the most important difference between the bee samples of the Thrace region was seen in the wing veinal angles.

The results of the tRNA^{leu}-cox2 sequencing of the Thrace honeybees

The analysis of the sequence data at the tRNA^{leu}-cox2 region produced twenty-two different haplotypes, of which sixteen were novel), and all of them were ascribable to the East European C-lineage and characterized by the presence of a single Q sequence corresponding to the predicted composition of this region. The mtDNA fragment corresponded to the positions 3363-3935 bp (Crozier & Crozier, 1993). The polymorphic sites of the intergenic tRNA^{leu}-cox2 region of the previous C2 haplotypes and the haplotypes that are found in this study, GenBank

accession numbers, and the variations with the nucleotide positions were summarized in Tab. 4. Sixteen polymorphic sites were detected within the sequenced 571-573 bp mtDNA fragment, and the sequences were deposited in the GenBank database with accession numbers MH939332- MH939353 (Tab. 4). In our study, 104 samples out of 217 (~48%) were found to be exactly the same as the published C2d haplotype in the NCBI Genbank database (accession numbers FJ824584, FJ824585, FJ037777, JQ977701, JF723977). C2d was found as a common haplotype all over the Thrace region, especially in Tekirdağ (0.78), Kırklareli (0.24), Edirne (0.67) and Gökçeada Island (0.50) (Tab. 4 and Tab. 5). In this study, the C2d sequence was deposited to the GenBank database with the accession number MH939332. Along with the C2d haplotype, previously published C2s, C2v, C2i, C2j, and C2h haplotypes were also observed in the Thrace region less frequently, forty-four samples out of 217 (20.27%). The DNA sequences of the *A. m. carnica*, *A. m. macedonica*, and *A. m. caucasica* reference samples were found exactly the same as the published C2v, C2d and C2h haplotypes, respectively. The remaining sixty-nine samples from different parts of Thrace, Çanakkale, and Gökçeada Island

Table 4.

The polymorphic sites of the intergenic tRNA^{leu}-cox2 region of the previous C2 haplotypes and the haplotypes that are found in this study, NCBI GenBank accession numbers and the citing references are given. The mtDNA fragment corresponds to the positions 3363-3935 bp published by Crozier & Crozier (1993).

Haplotypes	bp	3406	3410*	3424*	3425*	3428 ¹	3442	3449	3474	3488*	3512*	3514	3535	3536*	3567	3569	3573*	3575	3576*	3588	3589	3605	3634	3664*	3670	3735	3769	3771	References
JQ977700_C2c (<i>A. m. carnica</i>)	572	T	A	A	T	-	T	T	T	A	T	T	A	A	A	A	A	-	C	G	-	.	T	T	T	T	C	C	Sušnik et al., 2004; Muñoz et al., 2009, 2012
JQ977701 / FJ824585_C2d (<i>A. m. carnica</i> / <i>A. m. macedonica</i>)	572/ 526	.	.	.	.	-	.	.	.	.	.	.	.	.	.	.	.	-	.	.	-	.	C	.	.	.	T	.	Sušnik et al., 2004; Muñoz et al., 2009, 2012
JQ977702_C2e (<i>A. m. carnica</i>)	571	.	.	.	.	-	.	.	.	.	.	.	.	.	.	.	.	-	.	.	-	.	C	.	.	.	T	.	Kozmus et al., 2007; Muñoz et al., 2009
FJ357806_C2f (<i>A. m. medja</i>)	571	.	.	.	.	-	.	.	.	.	.	.	.	.	.	.	.	-	.	.	-	.	C	.	.	.	T	.	Özdil et al., 2009
FJ357807_C2g (<i>A. m. medja</i>)	571	.	.	.	.	-	.	.	.	.	.	.	.	.	T	.	.	-	.	.	-	.	C	.	.	.	T	.	Özdil et al., 2009
FJ357808_C2h (<i>A. m. caucasica</i>)	514	.	.	.	.	-	.	.	.	.	.	.	.	.	T	.	.	-	.	.	-	.	C	.	.	.	T	T	Özdil et al., 2009
FJ447491_C2i (<i>A. m. carnica</i>)	517	.	.	.	.	-	.	.	.	.	.	.	.	.	.	.	.	-	.	A	-	.	C	.	.	.	T	.	Nedić et al., 2009
JF723978_C2j (<i>A. m. carnica</i>)	572	.	.	.	.	-	.	.	.	.	.	.	.	.	T	.	.	-	.	.	-	.	C	.	.	.	T	.	Nedić et al., 2009; Corioian et al., 2014
GQ433624_C2k (<i>A. m. carnica</i>)	401	.	.	.	.	-	.	.	.	.	.	.	.	.	.	.	.	-	.	A	-	.	C	.	.	.	T	.	Razpet et al., 2009 unpub- lished
GQ433625_C2l (<i>A. m. carnica</i>)	401	.	.	.	.	-	.	.	.	.	.	.	.	.	.	.	.	-	.	.	-	.	.	.	.	.	.	.	Razpet et al., 2009 unpub- lished
GQ433626_C2m (<i>A. m. carnica</i>)	402	C	.	.	.	-	.	.	.	.	.	.	.	.	.	.	.	-	.	.	-	.	.	.	.	.	.	.	Razpet et al., 2009 unpub- lished
GQ433627_C2n (<i>A. m. carnica</i>)	402	.	.	.	.	-	.	.	.	.	.	.	.	.	.	.	.	-	.	.	-	.	.	.	.	.	.	.	Razpet et al., 2009 unpub- lished
JQ977704_C2o (<i>A. m. carnica</i>)	571	.	.	.	.	-	.	.	.	.	.	A	.	.	.	.	.	-	.	.	-	.	C	.	.	.	T	.	Muñoz et al., 2012
JQ977705_C2p (<i>A. m. carnica</i>)	571	.	.	.	.	-	.	.	.	.	.	.	.	.	.	.	.	-	.	.	-	.	C	.	.	.	T	.	Muñoz et al., 2012; Corioian et al., 2014
HM117905_C2q (<i>A. m. carnica</i>)	549	.	.	.	.	-	.	.	.	.	.	.	.	.	.	.	.	-	.	.	-	.	C	.	.	.	T	.	Muñoz 2013/ Corioian et al., 2014
HM117906_C2r (<i>A. m. carnica</i>)	546	.	.	.	.	-	.	.	.	.	.	.	T	.	.	.	.	-	.	.	-	.	C	.	.	.	T	.	Corioian et al., 2014
JF723979_C2s JQ973663_C2t JQ973664_C2v	572 572 572	.	.	.	.	-	.	.	.	.	.	A	.	.	T	.	.	-	.	.	-	.	C	.	.	.	T	.	Muñoz 2013 Muñoz 2013 Muñoz 2013
JQ754649_C2x	573	.	.	.	.	-	.	.	.	.	.	.	.	.	.	.	.	-	A	.	-	.	C	.	.	.	T	.	Corioian et al., 2014
JQ754650_C2y	571	.	.	.	.	-	.	.	.	.	.	.	.	.	.	.	.	-	.	.	-	.	.	.	.	.	.	.	Muñoz 2013; Corioian et al., 2014
JQ754648_C2z	572	.	.	.	.	-	.	.	.	.	.	.	.	.	.	.	.	-	.	.	-	.	.	.	.	.	.	.	Muñoz 2013/ Corioian et al., 2014

* The positions that are indicated with asterics are novel polymorphisms, character “-” shows deletions found.

Table 4 continuing

The haplotypes found in this study are given below. Novel haplotypes are written in bold.
The designations of the new haplotypes follow those of Franck et al. (2000).

MH939332_ C2d	572	.	.	.	.	-	.	.	.	.	.	.	.	.	.	-	.	.	-	.	C	.	.	.	T	.
MH939333_ C2d1	571	.	.	.	.	-	.	.	.	.	.	.	.	-	.	.	.	-	.	.	C	.	.	.	T	.
MH939334_C2s	572	.	.	.	.	-	.	.	.	.	A	.	.	.	.	-	.	.	-	.	C	.	.	.	T	.
MH939335_C2v	572	.	.	.	.	-	.	.	.	.	.	.	.	.	.	-	.	.	-	.	.	.	.	.	T	.
MH939336_ C2v1	572	.	.	.	.	-	.	.	.	.	.	.	.	.	.	-	.	A	-	.	.	.	.	.	T	.
MH939337_ C2v2	572	.	.	.	.	-	.	.	.	.	.	.	.	.	.	-	T	.	-	.	.	.	.	.	T	.
MH939338_ C2d2	572	.	.	.	.	-	.	.	.	.	C	.	.	.	.	-	.	.	-	.	C	.	.	.	T	.
MH939339_C2i	572	.	.	.	.	-	.	.	.	.	.	.	.	.	.	-	.	A	-	.	C	.	.	.	T	.
MH939340_ C2i2	572	.	T	.	.	-	.	.	.	.	.	.	.	.	.	-	.	.	-	.	C	.	.	.	T	.
MH939341_ C2i3	571	.	T	-	.	-	.	.	.	.	.	.	.	.	.	-	.	.	-	.	C	.	.	.	T	.
MH939342_ C2s1	572	.	T	.	.	-	.	.	.	.	A	.	.	.	.	-	.	.	-	.	C	.	.	.	T	.
MH939343_ C2c1	572	.	T	.	.	-	.	.	.	.	.	.	.	.	.	-	.	.	-	.	.	.	.	.	.	.
MH939344_ C2i1	572	.	T	.	.	-	.	.	.	.	.	.	.	.	.	-	.	A	-	.	C	.	.	.	T	.
MH939345_C2j	572	.	.	.	.	-	.	.	.	.	.	.	T	.	.	-	.	.	-	.	C	.	.	.	T	.
MH939346_C2h	572	.	.	.	.	-	.	.	.	.	.	.	T	.	.	-	.	.	-	.	C	.	.	.	T	T
MH939347_ C2d3	572	.	.	.	C	-	.	.	.	.	.	.	.	.	.	-	.	.	-	.	C	.	.	.	T	.
MH939348_ C2d4	572	.	.	.	.	-	.	.	.	.	.	.	.	.	.	T	-	.	.	-	C	.	.	.	T	.
MH939349_ C2h1	572	.	.	.	.	-	.	.	.	.	.	.	.	.	.	-	.	.	-	.	C	.	.	.	T	T
MH939350_ C2h2	573	.	.	.	.	-	.	.	.	.	.	.	.	.	.	-	.	.	G	.	C	.	.	.	T	T
MH939351_ C2h3	571	.	.	.	.	-	.	.	.	.	.	.	.	.	.	-	.	.	-	.	C	.	.	.	T	T
MH939352_ C2s2	572	.	.	.	.	-	.	.	.	.	A	.	.	.	.	-	.	.	-	.	C	C	.	.	T	.
MH939353_ C2s3	571	.	.	.	.	-	.	.	.	.	.	.	.	.	.	-	.	.	-	.	C	C	.	.	T	.

revealed sixteen novel haplotypes. Based on the results from this survey, existing haplotype names were revised and updated following a nomenclature system established earlier (C1-C2 system) and extended herein for the intergenic region. Novel haplotypes, C2d2, C2v1, and

C2v2, were observed in Tekirdağ and Kırklareli provinces, while C2i1, C2i2, C2i3, C2s1 and C2c1 haplotypes were only observed in the Kırklareli province, especially along the Istranca Mountain ridges and higher regions (Tab. 5). Along with these five new haplotypes found in Kırklareli,

Table 5.
Number of analyzed colonies (n), haplotype
distributions in the honey bee colonies from
Thrace and the reference populations

Location	n	C2d	C2d1	C2s	C2v	C2v1	C2v2	C2d2	C2i	C2i1	C2i2	C2i3	C2s1	C2c1	C2j	C2h	C2d3	C2d4	C2h1	C2h2	C2h3	C2s2	C2s3
Tekirdağ	50	39		2	5	1	2	1															
Kırklareli	67	16		1	8	2		1	2	3	22	6	2	3	1								
Edirne	49	33	1	2	5				4							2	2						
Çanakkale	31	6	8	3											5			2	2	2	3		
Gökçeada	20	10	1	4													1					1	3
	217	104	10	12	18	3	2	2	6	3	22	6	2	3	6	2	2	3	2	2	3	1	3
<i>A. m. carnica</i>	2				2																		
<i>A. m. macedonica</i>	2	2																					
<i>A. m. caucasica</i>	2															2							

twelve haplotypes out of 22 were observed in this region near the Istranca Mountains. On the other hand, C2h, C2h1, C2h2, C2h3, and C2d3 C2d4 haplotypes were also found only

in the Edirne and Çanakkale (Asian/Anatolian side) provinces, and these haplotypes were found to be similar to the previously published *A. m. caucasica* haplotypes (Özdil et al., 2009; Solorzano et al., 2009). Finally, the novel C2s2 and C2s3 haplotypes were only found in the Gökçeada Island populations (Tab. 5).

The phylogeny of the haplotypes based on the tRNA^{leu}-cox2 intergenic region in this study with the published Genbank reference haplotypes revealed mainly two, indeed three different clusters (Fig 3); C2c, C2l, C2m, C2n C2v, C2y, C2z and novel haplotypes found in this study C2c1, C2v1, C2v2 were clustered together. These novel haplotypes were only observed in the European part of Turkey-Thrace region. The second cluster was comprised of C2t, C2j, C2g, and C2h and novel C2h1, C2h2, C2h3 haplotypes, which had been previously defined as *caucasica*-like haplotypes. The remaining published haplotypes (C2d, C2k, C2l, C2p, C2q, C2o, C2s, C2r. etc.) and newly found C2d1-C2d4, C2s1-C2s3, C2i1-C2i3 haplotypes were clustered together in the third cluster (Fig. 3).

In this study, a NeighborNet network was constructed by the SplitsTree4 (version 4.1) program with the published and newly found haplotypes (Fig. 4). According to the NeighborNet network, all haplotypes were split into three main groups, which had been found in agreement with the morphometric and phylogenetic studies.

DISCUSSION

In the present study, we wanted to reveal the morphometric and genetic structure of the honey bees found in the European Part of Turkey, the Thrace region, located in the southeast of the Balkan Peninsula. Previous morphometric studies by Ruttner (1988), classified Thrace honey bees of Turkey as *A. m. anatoliaca*, but afterwards morphometric studies (Güler & Kaftanoğlu, 1999; Güler et al., 2010), mtDNA studies (Smith et al., 1997; Palmer et al., 2000; Kandemir et al., 2006; Ünal & Özdil, 2018) and microsatellites (Bodur et al., 2007) revealed that Thrace honey bees were different from *A. m. anatoliaca*. Not long ago, the Thrace




Fig. 3. Neighbor-joining tree based on intergenic tRNA^{leu}-cox2 region sequences performed on nucleotide distances computed by Kimura's (1980) formula through MEGA6 (Tamura et al., 2013).

■ Indicates novel haplotypes that were found in this study.

honey bee populations were considered as a distinct ecotype of *A. m. carnica* by the Republic of Turkey, Ministry of Agriculture and Forestry (Anonymous, 2020). The populations of the Thrace region (Tekirdağ, Edirne, and Kırklareli) were found similar to one another in relation to the morphological structure. However, the level of morphological similarity (8%) was found very low in this study, because the region had received uncontrolled queen bee introduction from different subspecies including the Caucasian or Anatolian honey bees. Güler (2010) had reported that the genetic mixture of a native bee population of a region was significantly affected by the use of external queen bees.

According to previous morphological and molecular studies, the honey bee populations of the Thrace region can be considered as an ecotype of *Apis mellifera carnica* (Palmer et al., 2000; Bodur et al., 2007; Güler et al., 2010; Ünal & Özdiş, 2018). The Carniolan honey bees have been reported to have important distinguishing morphological characters (Kauhausen-Keller et al., 1997; Ruttner, 1988; Güler et al., 2010). For example, according to Ruttner (1988), the most important distinctive morphological character of the Carniolan subspecies was the cubital index (CI; 2.589), while Dawino protocol and Güler et al. (2010) reported that the most important distinguishing character was the wing B₄ veinal angle (110° and 106°). In this study, the worker




Fig. 4. A NeighborNet Network constructed from SplitsTree4 (version 4.1)
 **Novel haplotypes found in this study are written in blue.

bee samples of the Thrace region have shown similarity to Carniolan subspecies in terms of the cubital index, B_4 veinal angle and some of the other characters (A_4 , D_7 , J_{10} , J_{16} and N_{23}) (Ruttner, 1988; Güler & Kaftanoğlu, 1999; Güler et al., 2010; Kulıci & Kume, 2014)

Although Ruttner (1988) stated that the Thrace honey bee populations were morphologically similar to the *Apis mellifera anatoliaca*, we found that the Thrace honey bees had lower similarity with the Anatolian subspecies. For example, the average scutellum colour of the Anatolian subspecies been found to be 5.83 scales (Güler & Kaftanoğlu, 1999), whereas in this study, the average scutellum color was found 0.543 in the Thrace honey bee populations.

Due to the ecological differences (flora, climatic, etc.) in the Balkan Peninsula, different ecotypes of the Carniolan subspecies have been originated, and the Thrace honey bees can be one of them because of the high variation that was observed between the populations of *A. m. carnica* in their original territory (Moritz, 1991; Poklukar & Kezic, 1994; Lodesani & Costa, 2003; Kulići & Kume, 2014). Indeed, according to Ruttner (1988), the Carniolan subspecies was

reported to have the most strains in the world in its natural geography.

On the other hand, the Gökçeada Island samples were clustered in a narrow space in the coordinate system (Fig. 2). The most important reason for that, the island is a completely isolated area and it maintains its natural position. The island worker bee samples were found to be completely different from the others in terms of seventeen characters. For example, the island bee has the longest hair (HL) and the largest body size (T_3+T_4) structures, which showed that its populations maintained their morphological structure compared to the previous studies (Güler & Kaftanoğlu, 1999). It is emphasized that the island's cold and continuous winds were considered to be the main reason for these morphological differences.

In this study, we also provide the most comprehensive survey of DNA sequencing of the tRNA^{leu}-cox2 region whichever reported for the European part of Turkey-Thrace region, Gökçeada Island and Çanakkale province (Gallipoli Peninsula and west Anatolian side). The C1/C2 haplotypes of the tRNA^{leu}-cox2 region were first reported by Franck et al., (2000) and the only difference

between the C1 and C2 was a single C nucleotide deletion (h) at the 3428th position of the mitochondrial genome. Franck et al. (2000) reported the C1, C2a and C2b haplotypes in their study, and the C2c and C2d haplotypes were observed by Sušnik et al. (2004) in Slovenia, Croatia (*A. m. carnica*) and Greece (*A. m. macedonica*), respectively. Also, the C2d haplotype was reported in all of the studied honey bees from Macedonia, and morphometrical analyses confirmed that those workers with the C2d haplotype were similar to *A. m. macedonica*, whereas the workers with the C2c haplotype were similar to *A. m. carnica* (Muñoz et al., 2009). Although most of the studies attributed the C2d haplotype to *A. m. macedonica*, subsequent studies have revealed that the C2d haplotype was also found in *A. m. carnica* (Kozmus et al., 2007; Muñoz et al., 2009, 2012; Coroian et al., 2014). After the C2c/C2d classification of the following haplotypes pertaining to the Balkan Peninsula, the novel haplotypes have been described but less frequently. The C2e haplotype was first reported in honey bees from Serbia (*A. m. carnica*) (Kozmus et al., 2007), and the C2h haplotype was first reported in *A. m. caucasica* samples from Turkey (Özdil et al., 2009). Another haplotype, the C2i, was first reported in Ikaria/ Greece (Muñoz et al., 2009) and later from Serbia (Nedić et al., 2009; Muñoz et al., 2012). The C2j haplotype was first reported in honey bees from Serbia (Nedić et al., 2009) and later in Romania (Coroian et al., 2014). The DNA sequences of the C2k, C2l, C2m and C2n haplotypes of the tRNA^{leu}-cox2 region were deposited to the NCBI GenBank database (GQ433624-GQ433627) but were unpublished. The haplotypes of the C2o (JQ977704) and C2p (JQ977705) were reported in Serbia (Muñoz et al., 2012). The additional haplotypes C2q, C2r, C2t, C2v, C2x, C2y and C2z were also reported in honey bees from Romania, and the C2s haplotype (JF723979) was first reported in honey bees from Hungary (Muñoz et al., 2012; Coroian et al., 2014). In our study, C2d was found to be the common haplotype (~48%) in the Thrace region. Previously published C2s, C2v, C2i, C2j and C2h haplotypes were also observed in the Thrace

region but less frequently (20.27%). Additionally, sixteen novel haplotypes were also obtained. Coroian et al. (2014) reported that according to the morphometrical analyses, the workers with the C2c haplotype were more similar to *A. m. carnica* and confirmed the haplotypes C2c, C2v, C2y and C2z were *carnica*-like haplotypes, whereas the C2d, C2e, C2f, C2p, C2q, C2r, C2s, C25-C28 were *macedonica*-like haplotypes. In our study, a neighbor-joining tree based on the intergenic tRNA^{leu}-cox2 region sequences was drawn and three different clusters were obtained. The first cluster mainly consisted of the *A. m. carnica* haplotypes (C2c, C2v) and newly found C2v1-Cv2, C2c2 that were found in the Thrace region, while the second cluster consisted of previously defined *caucasica*-like haplotypes. The remaining published haplotypes (C2d, C2k, C2l etc.) and newly found C2d1-C2d4, C2s1-C2s3 and C2i1-C2i3 haplotypes were clustered together in the third cluster which may be defined as *macedonica*-like haplotypes. According to the network, *carnica*-like haplotypes, *macedonica*-like haplotypes and *caucasica*-like haplotypes were again split from each other. When we consider the geographical distribution of the haplotypes, *caucasica*-like haplotypes were mainly found in the Asian/Anatolian side of Çanakkale and Edirne provinces, and this may be due to the commercial queen rearing, especially Caucasian queen import to this region. *Macedonica*-like haplotypes were mainly observed in honey bees from the European part of Turkey-Thrace region (Tekirdağ, Kırklareli and Edirne) and partly from Gökçeada Island. *Carnica*-like haplotypes were only found in honey bees from Kırklareli and especially the Istranca Mountain ridges and higher regions.

The genetic origin of honey bees found in the European part of the Turkey-Thrace region has been a mystery for a long time. First, Thrace honey bees were classified as *A. m. anatoliaca*, but new molecular and morphometric studies have revealed a different honey bee subspecies/ecotype in this region which showed similar morphometric and genetic characteristics to *A. m. macedonica* and *A. m. carnica*. According to the results of this study, the Thrace honey bees of

Turkey had lower similarity with *A. m. anatoliaca* or *A. m. caucasica*, but caucasian queen import may have a negative effect on the conservation of the genetic resources of the Thrace region. It is obvious that the Marmara Sea and the Gallipoli Peninsula also served as a natural barrier for the propagation of the *A. mellifera* subspecies. On the other hand, this study found that Thrace honey bees were more similar to *A. m. macedonica* through mtDNA sequence analysis, and that especially *carnica*-like honey bees were only found near the Istranca mountain ridges in the Kırklareli province while *macedonica*-like honey bees were observed all around the Thrace region in the European part of Turkey. Recently, the Republic of Turkey, Ministry of Agriculture and Forestry, have registered Thrace honey bees as an ecotype of *A. m. carnica*, and according to our results some of the Thrace honey bee populations may be both *A. m. carnica* and *A. m. macedonica* but the assignment to the latter subspecies seems more likely due to its geographic range. In conclusion, it is very important for these genetic resources and ecotypes to be protected in their original habitats and conservation strategies should be performed for the future.

ACKNOWLEDGMENTS

The authors are deeply indebted to numerous people that have contributed to this study for providing honey bee reference samples; Ljubiša Ž. Stanisavljević for providing *A. m. carnica* samples from Serbia and Leonidas Charistos for providing *A. m. macedonica* samples from Greece. Financial support for this research was provided by The Scientific and Technological Research Council of Turkey-TUBITAK through the Project 3001-TOVAG 1140883, Project Coordinator Fulya Özdil.

AUTHORS CONTRIBUTION

FO conceived this research, submitted the project and designed the experiments; DO and FO determined the sampling locations and participated in the sampling of the honey bees; AG participated in the design and interpretation of

the morphometric data; SY and AA performed the molecular and morphometric experiments and analysis; FO, RI and AG wrote the paper and participated in its revisions. All authors read and approved the final manuscript.

CONFLICT OF INTEREST

The authors declare that they have no conflict of interest.

REFERENCES

- Anonymous (2020). Resmi Gazete 2020/33, 05.12.2020 <https://www.resmigazete.gov.tr/eskiler/2020/12/20201205-5.htm>
- Bodur, Ç., Kence, M., Kence, A. (2007). Genetic structure of honeybee, *Apis mellifera* L. (Hymenoptera: Apidae) populations of Turkey inferred from microsatellite analysis. *Journal of Apicultural Research*, 46(1), 50-56. <https://doi.org/10.1080/00218839.2007.11101366>
- Bouga, M., Alaux, C., Bienkowska, M., Büchler, R., Carreck, N. L., Cauia, E., ... Wilde, J. (2011). A review of methods for discrimination of honey bee populations as applied to European beekeeping. *Journal of Apicultural Research*, 50(1), 51-84. <https://doi.org/10.3896/IBRA.1.50.1.06>
- Bouga, M., Harizanis, P. C., Kilias, G., Alahiotis, S. (2005). Genetic divergence and phylogenetic relationships of honey bee *Apis mellifera* (Hymenoptera: Apidae) populations from Greece and Cyprus using PCR - RFLP analysis of three mtDNA segments. *Apidologie*, 36(3), 335-344. <https://doi.org/10.1051/apido:2005021>
- Chen, C., Liu, Z., Pan, Q., Chen, X., Wang, H., Guo, H., ... Shi, W. (2016). Genomic Analyses Reveal Demographic History and Temperate Adaptation of the Newly Discovered Honey Bee Subspecies *Apis mellifera* sinixinyuan n. ssp. *Molecular Biology and Evolution*, 33(5), 1337-1348. <https://doi.org/10.1093/molbev/msw017>
- Cooley, W. W., & Lohnes, R. R. (1971). *Multivariate Data Analysis*. John Wiley and Sons Inc., New York 1971,

364 S. <https://doi.org/10.1002/bimj.19730150413>

Coroian, C. O., Muñoz, I., Schlüns, E. A., Paniti-Teleky, O. R., Erler, S., Furdui, E. M., ... Moritz, R. F. A. (2014). Climate rather than geography separates two European honeybee subspecies. *Molecular Ecology*, 23(9), 2353-2361. <https://doi.org/10.1111/mec.12731>

Crozier, R. H., & Crozier, Y. C. (1993). The mitochondrial genome of the honeybee *Apis mellifera*. Complete sequence and genome organization. *Genetics*, 133(1), 97-117.

Franck, P., Garnery, L., Loiseau, A., Oldroyd, B. P., Hepburn, H. R., Solignac, M., Cornuet, J. M. (2001). Genetic diversity of the honeybee in Africa: Microsatellite and mitochondrial data. *Heredity*, 86, 420-430. <https://doi.org/10.1046/j.1365-2540.2001.00842.x>

Franck, P., Garnery, L., Solignac, M., Cornuet, J. M. (2000). Molecular confirmation of a fourth lineage in honeybees from the Near East. *Apidologie*, 31(2), 167-180. <https://doi.org/10.1051/apido:2000114>

Garnery, L., Cornuet, J. M., Solignac, M. (1992). Evolutionary history of the honey bee *Apis mellifera* inferred from mitochondrial DNA analysis. *Molecular Ecology*, 1(3), 145-154. <https://doi.org/10.1111/j.1365-294X.1992.tb00170.x>

Güler, A. (2010). A morphometric model for determining the effect of commercial queen bee usage on the native honeybee (*Apis mellifera* L.) population in a Turkish province. *Apidologie*, 41(6), 622-635. <https://doi.org/10.1051/apido/2010037>

Güler, A., Bek, Y., Guven, H. (2010). The Importance of Morphometric Geometry on Discrimination of Carniolan (*Apis mellifera carnica*) and Caucasian (*A. m. caucasica*) Honey Bee Subspecies and in Determining Their Relationship to Thrace Region Bee Genotype. *Journal of the Kansas Entomological Society*, 83(2), 154-162.

Güler, A., & Kaftanoğlu, O. (1999). Morphological Characters of Some Important Races and Ecotypes of Turkish Honeybees (*Apis mellifera* L.)-I. *Turkish*

Journal of Veterinary and Animal Sciences, 23(Suppl.3), 565-570.

Hall, T. A. (1999). BIOEDIT: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/ NT. *Nucleic Acids Symposium Series*.

Han, F., Wallberg, A., Webster, M. T. (2012). From where did the western honeybee (*Apis mellifera*) originate? *Ecology and Evolution*, 2(8), 1949-1957. <https://doi.org/10.1002/ece3.312>

Huson, D. H., & Bryant, D. (2006). Application of phylogenetic networks in evolutionary studies. *Molecular Biology and Evolution*, 23(2), 254-267. <https://doi.org/10.1093/molbev/msj030>

Kandemir, I., Kence, M., Sheppard, W. S., Kence, A. (2006). Mitochondrial DNA variation in honey bee (*Apis mellifera* L.) populations from Turkey. *Journal of Apicultural Research*, 45(1), 33-38. <https://doi.org/10.1080/00218839.2006.11101310>

Kauhausen-Keller, D., Ruttner, F., Keller, R. (1997). Morphometric studies on the microtaxonomy of the species *Apis mellifera* L. *Apidologie*, 28(5), 295-307. <https://doi.org/10.1051/apido:19970506>

Kekecoglu, M., Bouga, M., Soysal, M. I., Harizanis, P. (2009). Genetic divergence and phylogenetic relationships of honey bee populations from Turkey using PCR-RFLP'S analysis of two mtDNA segments. *Bulgarian Journal of Agricultural Science*, 15(6), 589-597.

Kozmus, P., Stevanović, J., Stanimirović, Z., Stojić, V., Kulišić, Z., Meglič, V. (2007). Analysis of mitochondrial DNA in honey bees (*Apis mellifera*) from Serbia. *Acta Veterinaria*, 57(5-6), 465-476. <https://doi.org/10.2298/AVB0706465K>

Kuliçi, M., & Kume, K. (2014). Study on morphometric traits of the Albanian bees. *Albanian Journal of Agricultural Science, Special Issue*, 419-425.

Lodesani, M., & Costa, C. (2003). Bee breeding and genetics in Europe. *Bee World*, 84(2), 69-85. <https://doi.org/10.1007/s12061-003-0001-1>

doi.org/10.1080/0005772X.2003.11099579

Martimianakis, S., Klossa-Kilia, E., Bouga, M., Kiliyas, G. (2011). Phylogenetic relationships of Greek *Apis mellifera* subspecies based on sequencing of mtDNA segments (COI and ND5). *Journal of Apicultural Research*, 50(1), 42-50. <https://doi.org/10.3896/IBRA.1.50.1.05>

Meixner, M. D., Leta, M. A., Koeniger, N., Fuchs, S. (2011). The honey bees of Ethiopia represent a new subspecies of *Apis mellifera*-*Apis mellifera simensis* n. ssp. *Apidologie*, 42(3), 425-437. <https://doi.org/10.1007/s13592-011-0007-y>

Meixner, M. D., Costa, C., Francis, R. M., Uzunov, A., Kryger, P., Carreck, N. L., Hatjina, F., Büchler, R. (2014). Honey bee genotypes and the environment. *Journal of Apicultural Research*, 53(2), 183-187. <https://doi.org/10.3896/ibra.1.53.2.01>

Meixner, M. D., Pinto, M. A., Bouga, M., Kryger, P., Ivanova, E., Fuchs, S. (2013). Standard methods for characterising subspecies and ecotypes of *Apis mellifera*. *Journal of Apicultural Research*, 52(4), 1-28. <https://doi.org/10.3896/IBRA.1.52.4.05>

Moritz, R. F. A. (1991). The limitations of biometric control on pure race breeding in *Apis mellifera*. *Journal of Apicultural Research*, 30(2), 54-59. <https://doi.org/10.1080/00218839.1991.11101234>

Muñoz, I., Dall'Olio, R., Lodesani, M., Rúa, P. D. La. (2009). Population genetic structure of coastal Croatian honeybees (*Apis mellifera carnica*). *Apidologie*, 40(6), 617-626. <https://doi.org/10.1051/apido/2009041>

Muñoz, I., Stevanovic, J., Stanimirovic, Z., De la Rúa, P. (2012). Genetic variation of *Apis mellifera* from Serbia inferred from mitochondrial analysis. *Journal of Apicultural Science*, 56(1), 59-69. <https://doi.org/10.2478/v10289-012-0007-9>

Nedić, N., Stanisavljević, L., Mladenović, M., Stanisavljević, J. (2009a). Molecular characterization of the honeybee *Apis mellifera carnica* in Serbia. *Archives of Biological Sciences*, 61(4), 587-598. <https://doi.org/10.2298/ABS0904587N>

Oleksa, A., Chybicki, I., Tofilski, A., Burczyk, J. (2011). Nuclear and mitochondrial patterns of introgression into native dark bees (*Apis mellifera mellifera*) in Poland. *Journal of Apicultural Research*, 50(2), 116-129. <https://doi.org/10.3896/ibra.1.50.2.03>

Özdil, F., Yıldız, M. A., Glenn Hall, H. (2009). Molecular characterization of Turkish honey bee populations (*Apis mellifera*) inferred from mitochondrial DNA RFLP and sequence results. *Apidologie*, 40(5), 570-576. <https://doi.org/10.1051/apido/2009032>

Palmer, M. R., Smith, D. R., Kaftanoğlu, O. (2000). Turkish honeybees: Genetic variation and evidence for a fourth lineage of *Apis mellifera* mtDNA. *Journal of Heredity*, 91(1), 42-46. <https://doi.org/10.1093/jhered/91.1.42>

Péntek-Zakar, E., Oleksa, A., Borowik, T., Kusza, S. (2015). Population structure of honey bees in the Carpathian Basin (Hungary) confirms introgression from surrounding subspecies. *Ecology and Evolution*, 5(23), 5456-5467. <https://doi.org/10.1002/ece3.1781>

Poklucar, J., & Kezic, N. (1994). Estimation of heritability of some characteristics of hind legs and wings of honeybee workers (*Apis mellifera carnica* Polm.) using the half-sibs method. *Apidologie*, 25(1), 3-11. <https://doi.org/10.1051/apido:19940101>

Ruttner, F. (1988). *Biogeography and Taxonomy of Honeybees*. https://doi.org/10.1007/978-3-642-72649-1_6

Ruttner, F., Tassencourt, L., Louveaux, J. (1978). Biometrical-statistical analysis of the geographic variability of *Apis mellifera* L. I. Material and Methods. *Apidologie*, 9(4), 363-381. <https://doi.org/10.1051/apido:19780408>

Sambrook, J., & Russel, D. W. (2001). Molecular Cloning, 3-Volume Set: A Laboratory Manual. In *Cold Spring Harbor Laboratory Press*.

Sheppard, W. S., Arias, M. C., Grech, A., Meixner, M. D. (1997). *Apis mellifera ruttneri*, a new honey bee subspecies from Malta. *Apidologie*, 28(5), 287-293.

<https://doi.org/10.1051/apido:19970505>

Sheppard, W. S., & Meixner, M. D. (2003). *Apis mellifera pomonella*, a new honey bee subspecies from Central Asia. *Apidologie*, 34(4), 367-375. <https://doi.org/10.1051/apido:2003037>

Smith, D. R., Slaymaker, A., Palmer, M., Kaftanoğlu, O. (1997). Turkish honey bees belong to the east Mediterranean mitochondrial lineage. *Apidologie*, 28(5), 269-274. <https://doi.org/10.1051/apido:19970503>

Solorzano, C. D., Szalanski, A. L., Kence, M., McKern, J. A., Austin, J. W., Kence, A. (2009). Phylogeography and population genetics of honey bees (*Apis mellifera*) from Turkey based on COI-COI sequence data. *Sociobiology*, 53(1), 237-246.

Spss. (2004). SPSS 13.0 Base User's Guide®. In *Text*. Sušnik, S., Kozmus, P., Poklukar, J., Meglič, V. (2004). Molecular characterisation of indigenous *Apis mellifera carnica* in Slovenia. *Apidologie*, 35(6), 623-636. <https://doi.org/10.1051/apido:2004061>

Tamura, K., Stecher, G., Peterson, D., Filipski, A., Kumar, S. (2013). MEGA6: Molecular evolutionary genetics analysis version 6.0. *Molecular Biology and Evolution*, 30(12), 2725-2729. <https://doi.org/10.1093/molbev/mst197>

Ünal, G., & Özdil, F. (2018). Genetic characterization of Thrace honey bee populations of Turkey: restriction and sequencing of inter cytochrome C oxidase I-II (CoxI-CoxII) genes. *Journal of Apicultural Research*, 57(2), 213-218. <https://doi.org/10.1080/00218839.2018.1426347>

Whitfield, C. W., Behura, S. K., Berlocher, S. H., Clark, A. G., Johnston, J. S., Sheppard, W. S., ... Tsutsui, W. D. (2006). Thrice out of Africa: Ancient and recent expansions of the honey bee, *Apis mellifera*. *Science*, 314(5799), 642-645. <https://doi.org/10.1126/science.1132772>