

Title: A recalcitrant case of facial verruca plana successfully treated with topical calcipotriol alone

Ömer Faruk Elmas¹, Mustafa Atasoy²; Necmettin Akdeniz³; Kemal Özyurt⁴

¹ MD, Ahi Evran University, Kırşehir, Turkey

² MD, Health Science University, Kayseri City Hospital, Kayseri, Turkey

³ MD, Istanbul Medeniyet University, Istanbul, Turkey

⁴MD, Ahi Evran University, Kırşehir, Turkey

Corresponding Author:

Ömer Faruk Elmas

Ahi Evran University, Dept of Dermatology, Kırşehir, 40000, Turkey

omerfarukmd@gmail.com

+905330260679

Keywords: calcipotriol, plane wart, verruca plana

Running title: Verruca plana treated with topical calcipotriol

The authors declare that they have no conflict of interest.

No funding source.

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1111/dth.13664

A recalcitrant case of facial verruca plana successfully treated with topical calcipotriol

Verruca plana is a benign epidermal proliferative cutaneous condition mainly caused by human papillomavirus (HPV) types 3 and 10. HPV 27, 28, 38, 41, and 49 may also rarely cause verruca plana¹. Face located plane warts usually cause important aesthetic concerns in patients and lead them to seek treatment. The management of verruca plana poses a therapeutic challenge for clinicians, as there is no single method that has been proven to be efficient for completely treating the disease.

An 18-year-old woman presented with an eight months history of asymptomatic numerous small brown lesions on the chin. She was otherwise healthy and her past medical history was unremarkable. The lesions did not respond to previously treatments including oral zinc, topical tretinoin and topical adapalene. Dermatological examination revealed numerous, plane, yellowish to brown irregular papular lesions ranging in size from 0.1 cm to 0.4 cm distributed on the left side of the chin (Figure 1a). Dermatoscopic examination showed numerous tiny dotted (pinpoint) vessels on a light brown background. No other lesions were observed elsewhere. Based on the clinical and dermatoscopic examination, a diagnosis of verruca plana was made. Daily simple application of topical calcipotriol ointment allowed a complete resolution after 8 weeks without adverse reactions (Figure 1b). No recurrence was observed during six months of follow up.

Common therapeutic options for verruca plana include cryotherapy, electrocoagulation, topical salicylic acid, topical retinoids, topical imiquimod, topical immunotherapy, laser or conventional/daylight photodynamic therapy (PDT)^{2,3}. Most of these options may cause dyspigmentation, pain, and even scarring. Furthermore, no treatment has been uniformly effective and, treatment resistance is common.

There are very few studies on the effectiveness of topical vitamin D derivatives in the treatment of viral warts. Imagawa et al reported a series of 17 cases of hand or foot located warts successfully treated with topical maxacalcitol⁴. Egawa et al, managed three immunocompromised cases with recalcitrant warts, using topical maxacalcitol with a half-day occlusive dressing technique⁵. Labandeira et al reported a case of giant wart successfully treated with topical calcipotriol⁶. Rind et al reported the successful use of topical calcipotriol in a case of anogenital wart in an infant⁷.

Recently, it has been shown that intralesional vitamin D3 injection may be a simple, safe and cost-effective treatment modality with minimal side effects. In a single-blinded placebo-controlled study investigating effectivity of intralesional vitamin D3 injection in the treatment of common warts, complete clearance of the target injected warts occurred in 40% of patients in cases group, while it occurred only in 5% of patients in control groups⁸.

Topical vitamin D3 derivatives are considered as first-line therapy in the treatment of chronic plaque psoriasis⁹. Biologic effects of vitamin D3 derivatives mainly include regulation of epidermal proliferation and modulation of cytokine production. Recent studies showed that vitamin D3 derivatives may also have an effect on cell death and angiogenesis. These biological actions may explain the efficacy of vitamin D3 derivatives for treating warts⁵.

Accepted Article

In conclusion, this report demonstrates that a simple application of topical calcipotriol may be an effective and safe treatment option for plane warts. However, the resolution of warts in one patient may not be proof of the efficacy of topical calcipotriol because plane warts may also resolve spontaneously in a certain subset of patients. Considering that the classical treatment options usually have limited efficiency and side effects such as pain, pigmentary changes and scarring, we would encourage placebo-controlled studies investigating the effectiveness and safety of topical vitamin D3 derivatives for treating plane warts.

Figure legends.

Figure1: **a.** Verruca plana. Numerous, plane, yellowish to brown, irregular papular lesions ranging in size from 0.1 cm to 0.4 cm distributed on the left side of the chin. **b.** Complete resolution of the lesions after daily application of topical calcipotriol for 8 weeks.

Figure 2: Dermoscopic examination showed numerous pinpoint vessels widely scattered over the light brown background.

References

1. Zhang K, Ren J, Ren JX, Liu C, Sun L. Successful treatment of verruca plana with NB-UVB: A case report. *Dermatol Ther.* 2019:e13207.
2. Zhang F, Shi L, Liu P, et al. A novel cosmetic and clinically practicable laser immunotherapy for facial verruca plana: Intense pulsed light combined with BCG-PSN. *Photodiagnosis Photodyn Ther.* 2018;22:86-90.
3. Borgia F, Coppola M, Giuffrida R, Cannavo SP. Excellent cosmetic result of daylight photodynamic therapy for facial flat warts in a child. *Photodiagnosis Photodyn Ther.* 2019;26:27-28.
4. Imagawa I, Suzuki H. Successful treatment of refractory warts with topical vitamin D3 derivative (maxacalcitol, 1alpha, 25-dihydroxy-22-oxacalcitriol) in 17 patients. *J Dermatol.* 2007;34(4):264-266.
5. Egawa K, Ono T. Topical vitamin D3 derivatives for recalcitrant warts in three immunocompromised patients. *Br J Dermatol.* 2004;150(2):374-376.

6. Labandeira J, Vazquez-Blanco M, Paredes C, Suarez-Penaranda JM, Toribio J. Efficacy of topical calcipotriol in the treatment of a giant viral wart. *Pediatr Dermatol*. 2005;22(4):375-376.
7. Rind T, Oiso N, Kawada A. Successful Treatment of Anogenital Wart with a Topical Vitamin D(3) Derivative in an Infant. *Case Rep Dermatol*. 2010;2(1):46-49.
8. Kareem IMA, Ibrahim IM, Mohammed SFF, Ahmed AA. Effectiveness of intralesional vitamin D3 injection in the treatment of common warts: Single-blinded placebo-controlled study. *Dermatol Ther*. 2019;32(3):e12882.
9. Mason J, Mason AR, Cork MJ. Topical preparations for the treatment of psoriasis: a systematic review. *Br J Dermatol*. 2002;146(3):351-364.



